

Contents

Unit 1 First Principles

Chapter 1: The Foundations of Scientific Inquiry 3

THE DIFFERENCE BETWEEN EVIDENCE AND TRUTH 4

Immanuel kant's *critique of pure reason* picks up where descartes leaves off 5

EMPIRICISM AND THE SCIENTIFIC METHOD 6

Observation 6

Question 7

Hypothesis 8

Experiment 10

Analysis 11

Conclusion 12

Report 12

AN EXAMPLE OF THE SCIENTIFIC METHOD IN ACTION 12

Observation 12

Question 13

Hypothesis 13

Experiment 13

FORWARD IN SCIENCE, FORWARD IN TRUTH 16

REFERENCES 16

FURTHER READING 16

Chapter 2: Introduction to Statistical Inference 17

DIFFERENT TYPES OF DATA 18

Every measurement is limited by imperfections in precision and accuracy 19

The 30–300 rule is used to determine the proper instrument of measure 20

The rules of significant figures define the precision of any measurement 22

FOR EVERY MEASURE THERE IS ERROR 22

Descriptive assessments of central tendency are used for accurate representations of the variability in a sample 24

“Normal” data can be plotted as a gaussian curve 27

The coefficient of variation can be used to measure variability 28

Data distributions that aren't normal often follow a poisson distribution 30

Nonnormal datasets can be normalized using data transformation 30

The central limit theorem can be employed to increase the predictability of your data distributions 31

INFERENTIAL STATISTICS: A BRIEF PRIMER 33

The probability level will define which results are statistically significant, and which are not 33

The use of confidence intervals and outliers helps in the analysis of statistical significance 34

The strength of our statistical analyses will depend heavily on our assumptions of the data 34

Independence 35

Single population 36

Normality 36

Homoscedasticity 36

Formulating a testable hypothesis is critical to both the scientific method and the use of inferential statistics 37

The most basic statistical tests are used to test for equality (or inequality) between two populations 38

CHOOSING THE RIGHT STATISTICAL METHOD 40

REFERENCES 42

FURTHER READING 42

Unit 2 Methods of Data Acquisition

Chapter 3: Experimental Design 45

STARTING WITH THE BASICS OF EXPERIMENTAL DESIGN 46

Method comparison 47

Single-element monitoring and assessment 48

Multiple-element correlation or causation 48

Complex synthetic analysis 49

Proper experimental design requires significant preparation 49

Literature review 49

Data requirements	50	Depth measures are used to extend spatial relationships into the vertical dimension	72
Research strategy	50	Time is required to define the succession of events	73
Scope of investigation	50	PHYSICAL VARIABLES FOR CONSIDERATION AND THE ESSENTIAL MEASURES OF OCEAN PHYSICS	75
Equipment and facilities needs	51	Mass is the most important way to quantify existence	75
Preliminary site survey(s)	51	Temperature affects the physical properties of all matter	76
Sampling effort	51	Hydrostatic pressure is used to describe the potential energy within a fluid	77
Sample size determination is the key to success in every research endeavor	53	Density describes the relationship between matter and the space it occupies	78
Dichotomous variables can be used to determine n_o	53	SUPPLEMENTAL MEASURES OF OCEAN PHYSICS	78
Continuous variables can also be used to determine n_o	54	Flow fields (and currents) are used to describe both the speed and direction of water movement	78
AN INTRODUCTION TO QUANTITATIVE SAMPLING METHODS	55	Wave geometry is used to discern wave patterns in a complex ocean	80
BASIC PLOT METHODS	56	Tides represent the influence of heavenly bodies on ocean dynamics	80
Sampling effort	56	Secchi depth is used as a simple way to define the optical properties of water	80
Plot shape	56	CHEMICAL VARIABLES	81
Plot size	58	Salinity is the fundamental measure of salt content in the ocean	82
Random sampling	59	Conductivity represents an indirect way to measure salinity using electricity	83
Simple random sampling is the foundation of most field methods	60	Metabolic nutrients are dissolved molecules which are necessary for life in the ocean	83
Stratified random sampling can be used to organize the study area into horizontal or vertical strata	60	SUPPLEMENTAL MEASURES OF OCEAN CHEMISTRY	84
Transect interval sampling is a variation of the belt/line transect method	61	Structural nutrients are dissolved molecules which can form solid mineral deposits	84
Adaptive sampling is an excellent method for detecting rare events	63	Dissolved O ₂ is one of the most influential molecules that affects the biogeochemistry of the ocean	84
BASIC PLOTLESS METHODS	63	Dissolved CO ₂ can be used to investigate gas exchange and ocean acidification dynamics	84
Plotless density estimators (PDEs) do not require exhaustive surveys of an area	63	BIOLOGICAL VARIABLES	85
Basic distance estimators (BDEs)	64	Species identification is the only way to describe the diversity of life in the ocean	85
Ordered distance estimators (ODEs)	64	Species abundance and biomass measures are used to quantify the amount of biologic matter	85
Angle-order estimators (AOEs)	64	Species biometrics represent those measures that suggest a specific biological function	86
Line transect density estimators (LTDEs) are the best PDE for “line-of-sight” search patterns	66	SUPPLEMENTAL MEASURES OF OCEAN BIOLOGY	87
Variable area transect (VAT)	66	Accessory pigments can be used to identify different species of plants and algae	87
The strong method (SM)	67	Organic C-N-P ratios define the nutritional status of the living biomass	87
Plotless methods for mobile subjects require a strategy different than most PDEs	67	Age structure and size class measures establish the demographics within a specific population	87
Roaming surveys	67		
Mark and recapture	68		
Catch per unit effort (CPUE)	68		
REFERENCES	68		
FURTHER READING	68		
Chapter 4: Oceanographic Variables	69		
ESTABLISHING THE FRAME OF REFERENCE	70		
DEFINING SPATIAL AND TEMPORAL BOUNDARIES	71		
Distance measures are a simple but critical way to define spatial relationships	72		

GEOLOGICAL VARIABLES	87	Titrimetric analysis quantifies hydrologic parameters by inducing a specific chemical reaction	105
Geomorphology and bathymetry describe the complex geometry at the ocean boundaries	88	Electronic analysis measures small changes in conductance to quantify hydrologic parameters	105
Suspended sediments provide the critical link between land and sea	88	REFERENCES	106
SUPPLEMENTAL MEASURES OF OCEAN GEOLOGY	89	FURTHER READING	106
Sediment grain characteristics provide forensic evidence of historical events	89	Chapter 6: Census Methods for Benthic Organisms	107
OTHER VARIABLES	89	CHOOSING THE RIGHT METHOD FOR EACH BIOTIC HABIT	108
Winds provide a strong influence on flow fields and surface currents	89	Starting simple is the key to every biological census	109
Barometric pressure can affect the hydrostatic pressure of ocean water beneath it	90	Take note whether the census includes open or closed populations	109
Air temperature can provide a delayed influence on water temperature	90	Benthic organisms require a two-dimensional census method	110
Precipitation will affect both the temperature and salinity of surface water in the ocean	90	CENSUS METHODS FOR THE SESSILE BENTHOS	111
REFERENCES	91	Rare species are best surveyed using the T-square plotless method	111
FURTHER READING	91	The distance sampling plot method is best for closed populations	112
Chapter 5: Common Hydrologic Census Methods	93	The line transect plot method is among the simplest to use	113
THE DIFFERENCE BETWEEN HYDROLOGY AND HYDROGRAPHY	94	True area surveys are accomplished using quadrat plot methods	114
WHEN TO MEASURE HYDROLOGIC DATA	94	Long-term studies may require a fixed quadrat plot method	116
Continuous sampling methods are best when measuring rapidly changing hydrologic variables	95	CENSUS METHODS FOR THE MOTILE BENTHOS	117
Underway systems provide continuous sampling during transit between multiple locations	95	Trapping and removal methods are the most common census strategy for motile benthic specimens	117
Moored systems provide continuous sampling at fixed locations only	96	Pitfall traps are best for capturing small, motile species above sea level	119
Discrete sampling methods are best when measuring slowly changing hydrologic variables	97	Attractant traps are excellent for capturing forager and scavenger species	119
WHERE TO COLLECT HYDROLOGIC DATA	99	Emergence traps are best used in the capture of cryptic species	120
Proper site selection is the key to success for any sampling strategy	99	Enclosure and exclosure traps are best suited for mesocosm studies	121
Fluvial study sites are those located along streams and rivers	100	Mark-and-recapture methods are a “trapless” solution to census motile species	121
Lacustrine and estuarine study sites are those located in bays and inland coastal areas	101	The pseudo-removal method can be used to estimate the size of a closed population by simply marking specimens	122
Marine study sites are those located along the coast or out to sea	102	The wileyto removal method for closed populations uses a combination of marked and removed specimens to assess population size	122
Don't forget the vertical dimension when planning your study sites	102	The multiple mark-and-recapture (schnabel) method for closed populations is among the most rigorous estimates of population size	123
HOW TO MEASURE HYDROLOGIC DATA	104	Open populations are more challenging to assess and require complex multiple mark-and-recapture methods	124
Colorimetric analysis uses color as a method to quantify hydrologic parameters	104		

REFERENCES 124

FURTHER READING 124

Chapter 7: Census Methods for Pelagic Organisms 125

LIFE IN A FLUID MEDIUM 126

Pelagic surveys require a definition of the vertical dimension 126

Size matters when choosing the proper gear and method for pelagic surveys 129

CENSUS METHODS FOR VIRUSES, BACTERIA, AND SMALL PHYTOPLANKTON 131

Virio- and bacterioplankton census methods require specialized equipment 132

Small (<10 μm) phytoplankton are most effectively collected by bottle 133

CENSUS METHODS FOR LARGE PHYTOPLANKTON, ZOOPLANKTON, AND LARVAE 133

The geometric dimensions of a plankton net determine its ideal function 133

Choosing the appropriate type of plankton tow requires some knowledge of the target organisms' vertical distribution in the ocean 135

The horizontal tow is best for sampling at a single, discrete depth layer 135

The vertical tow is best for integrated sampling of the entire water column 138

The double-oblique tow offers the best of both worlds, combining features of the horizontal and vertical tows 138

Plankton quantitation requires knowledge of the filtered volume of seawater 139

The sample volume from a plankton tow can be used to derive the population density of plankton in natural waters 141

Plankton Samples Must Be Properly Preserved for Later Analyses 141

The methods for processing plankton rely on representative subsamples 142

Counting and identifying plankton is a visual enterprise 143

CENSUS METHODS FOR FISHES AND OTHER NEKTON 144

Biometry statistics are critical to nekton surveys 146

The natural abundance of nekton can be estimated using "Catch Per Unit effort" (CPUE) data 147

NETS OR HOOK-AND-LINE CAPTURE METHODS FOR NEKTON 148

Seine nets can be easily deployed in shallow water by boat or by hand 148

Trawl nets are used in deeper waters and require the use of a vessel 149

Dredges must be pulled along the bottom to collect benthic organisms 150

Cast and push netting are best suited for use in bays, tidal creeks, and other small coastal areas 150

Gill netting is used to collect specimens from vast open-water areas, over extended periods of time 151

Trapping allows the collection of specimens that are unharmed during their capture 151

Hook and line methods are typically more selective than gill netting 151

How to define "effort" within CPUE analyses, based on the capture method 152

ANALYSIS OF PELAGIC COLLECTIONS 153

REFERENCES 153

FURTHER READING 153

Unit 3 Methods of Data Analysis

Chapter 8: Introduction to Univariate Analysis 157

THE FOUNDATIONS OF UNIVARIATE ANALYSIS 158

Univariate comparisons must be structured to possess only one scaled variable 159

COMMON STATISTICAL METHODS FOR UNIVARIATE ANALYSIS 162

The most common statistical tool for univariate analysis is the *t* test 163

The appropriate use of the *t* test depends on three critical assumptions 165

The single-sample *t* test is used to compare measurements against a historical or theoretical value 166

The paired-sample *t* test is used to test the dependence between two populations 166

Variance is used to determine whether to use the pooled-sample or welch's *t* test 167

The pooled-sample *t* test assumes equal variances between the populations being compared 168

Welch's *t* test is used when the variances of two populations are not equal 169

The one-way analysis of variance (ANOVA) is used to compare multiple populations simultaneously 169

COMMUNITY COMPARISONS USING UNIVARIATE ANALYSIS 171

Measures of species richness are the easiest way to compare communities 171

Simple community comparisons can be performed using ranked abundance data and the DAFOR/SACFOR methods 172

Biodiversity and its related measures are the cornerstones of community comparison methods 173

USING THE BEFORE-AFTER-CONTROL-IMPACT (BACI) APPROACH	177	USING CONCEPTUAL DIAGRAMS	206
MOVING BEYOND THE SIMPLICITY OF UNIVARIATE ANALYSIS	178	THE FUNDAMENTAL ELEMENTS OF MODELING	207
REFERENCES	179	Forcing functions are the fundamental equations that define the mathematical structure of the model	207
FURTHER READING	179	Variables are those elements within the model that can be quantified, and whose values are likely to change	208
Chapter 9: Introduction to Multivariate Analysis	181	Parameters and constants differ from variables in that they represent fixed quantities that do not change	209
INTRODUCTION TO CORRELATION ANALYSIS	182	FROM CONCEPT TO CREATION: WHAT NOW?	209
Correlation analysis can only be performed among continuous variables	182	REFERENCES	210
Correlation analysis can be used on no more than two dependent variables at a time	183	FURTHER READING	210
LINEAR REGRESSION ANALYSIS	184	Chapter 11: Model Structure	211
Correlations among normal data can be tested using pearson's correlation coefficient	185	DEFINING SPACE AND TIME	212
Correlations among nonnormal data can be tested using spearman's ranked correlation coefficient	187	Zero-dimensional models simulate what occurs at a single point	212
Correlations can also be used to make predictions using regression analysis	187	One-dimensional models simulate what occurs along a line	213
Regression analysis is used to test hypotheses and define confidence intervals within predictive mathematical models	189	Two-dimensional models simulate what occurs within a defined area	214
MULTIPLE REGRESSION AND CURVE ESTIMATION	191	Three-dimensional models simulate what occurs within a fixed volume	215
Multiple regressions that are nonlinear must be determined using curve estimation	194	Spatial dimensions must be resolved using a model grid	215
CORRELATION BETWEEN TWO DEPENDENT VARIABLES	195	CARTESIAN GRIDS	216
Canonical correlation (CC) is used to explore correlations between two sets of dependent variables	195	TELESCOPING GRIDS	217
Principal component analysis (PCA) is used to create two new sets of dependent variables and explore their correlation	196	CURVILINEAR GRIDS	218
FROM MEASUREMENTS TO MODELS	197	UNSTRUCTURED GRIDS	218
REFERENCES	198	VERTICAL GRIDS	218
FURTHER READING	198	The model timestep is used to resolve the temporal dimension	218
Unit 4 Methods of Data Assimilation (Modeling)		Striking a balance between model resolution and stability is paramount	220
Chapter 10: Fundamental Concepts in Modeling	201	BOUNDARY CONDITIONS	220
SO WHAT IS A MODEL?	202	COMPUTATIONAL SCHEMATA	222
MATHEMATICAL MODELS AS SCIENTIFIC TOOLS	204	Model currency is the unifying medium of exchange between variables within the model	223
		Mass and energy budgets must obey the law of conservation	223
		Model initialization requires that we begin with a "filled" model domain	224
		REFERENCES	229
		FURTHER READING	229

Chapter 12: Modeling Simple Dynamics 231

CONSTRUCTING YOUR FIRST MODEL 232
 The first rule of model construction: do your homework 232
 The second rule of model construction: follow your conceptual model 233
 The third rule of model construction: use the appropriate arrays 234
 The fourth rule of model construction: define the boundary conditions 235
 The fifth rule of model construction: establish your refuges 235
 The sixth rule of model construction: stay within your mass budget 236
 The unwritten rule of model construction: check your submodels for goodness of fit 237
EXAMPLES OF MODELING SIMPLE DYNAMICS 239
 Example 1: nitrogen dynamics of nitrification 239
 Example 2: nitrogen dynamics of excretion 241
 Example 3: nitrogen dynamics of production 243
REFERENCES 243
FURTHER READING 244

Chapter 13: Modeling Complex Dynamics 245

EMBRACING COMPLEXITY IN OUR MODEL(S) 246
 Complex processes are more easily linked by converting your conceptual models to conceptual equations 246
 A variable-oriented perspective will help you focus on what goes into (and what comes out of) each modeled process 246
 A process-oriented perspective will help you focus on which submodels should be included 247
 Conceptual equations are the best way to visualize how submodels should be connected with each other 248
EXAMPLES OF MODELING COMPLEX DYNAMICS 249
 Example 2 revisited: complex nitrogen dynamics of excretion 250
 Example 3 revisited: complex nitrogen dynamics of assimilation/production 256
MOVING ON TO BIGGER AND BETTER MODELS 260
REFERENCES 261
FURTHER READING 261

Chapter 14: Modeling Large System Dynamics 263

PUTTING IT ALL TOGETHER 264
TYPICAL HYDRODYNAMIC MODEL ELEMENTS 266
 Hydrodynamic models are solved by defining the forces at work within the spatial and temporal domain of fluid motion 266
 The unequal distribution of mass, as it relates to water density, creates pressure-gradient forces that influence water movement 268
 Frictional forces in the ocean play a significant role in water movement 269
 The influence of tides is as varied as it is powerful 269
TYPICAL BIOGEOCHEMICAL MODEL ELEMENTS 270
 The fundamentals of chemistry are the backbone of biogeochemical models 270
 Inorganic chemicals define the chemical nature of seawater, divorced from biology 270
 Organic chemicals in seawater are the calling cards of life 271
 Bacteria provide a fundamental link between the organic and inorganic chemistry of the sea 272
 Diagenesis represents the fusion of geology and chemistry 273
TYPICAL RADIATIVE MODEL ELEMENTS 273
 Light and heat must penetrate the atmosphere before they reach the ocean 273
 The radiation reaching the ocean surface is quickly attenuated as it travels through the water 275
TYPICAL ECOLOGICAL MODEL ELEMENTS 277
 Growth (∂g) describes the positive increment of biomass either from production among autotrophs or assimilation among heterotrophs 277
 Diffusive and advective transport (∂X) move biomass throughout the domain in the horizontal dimension 278
 Vertical migration and sinking (∂Z) move biomass throughout the domain in the vertical dimension 278
 Reproduction (∂Y) represents the fraction of growth that is invested in the next generation 278
 Respiration (∂R) is the loss of biomass attributed to gaseous wastes 278
 Excretion (∂E) is the loss of biomass attributed to liquid wastes 279
 Egestion (∂F) is the loss of biomass attributed to solid wastes 279
 Mortality (∂M) is always represented as a loss of biomass, but it comes in many forms 279

CLOSING THOUGHTS	280	Appendix C Common Algorithms for the Estimation of Important Water Properties	293
REFERENCES	280		
Appendices			
Appendix A Symbols and Abbreviations	281	Glossary	301
Appendix B Constants, Equivalents, and Common Units of Measure	287		