

Detailed Contents

PART I AN INTRODUCTION TO IMMUNO-BIOLOGY AND INNATE IMMUNITY

Chapter 1 Basic Concepts in Immunology 1

The origins of vertebrate immune cells. 2

Principles of innate immunity. 3

- 1-1 Commensal organisms cause little host damage while pathogens damage host tissues by a variety of mechanisms. 3
- 1-2 Anatomic and chemical barriers are the first defense against pathogens. 5
- 1-3 The immune system is activated by inflammatory inducers that indicate the presence of pathogens or tissue damage. 6
- 1-4 The myeloid lineage comprises most of the cells of the innate immune system. 7
- 1-5 Sensor cells express pattern recognition receptors that provide an initial discrimination between self and nonself. 8
- 1-6 Sensor cells induce an inflammatory response by producing mediators such as chemokines and cytokines. 9
- 1-7 Innate lymphocytes and natural killer cells are effector cells that share similarities with lymphoid lineages of the adaptive immune system. 11

Summary. 11

Principles of adaptive immunity. 11

- 1-8 The interaction of antigens with antigen receptors induces lymphocytes to acquire effector and memory activity. 12
- 1-9 Antibodies and T-cell receptors are composed of constant and variable regions that provide distinct functions. 13
- 1-10 Antibodies and T-cell receptors recognize antigens by fundamentally different mechanisms. 14
- 1-11 Antigen-receptor genes are assembled by somatic gene rearrangements of incomplete receptor gene segments. 15
- 1-12 Lymphocytes activated by antigen give rise to clones of antigen-specific effector cells that mediate adaptive immunity. 15
- 1-13 Lymphocytes with self-reactive receptors are normally eliminated during development or are functionally inactivated. 16
- 1-14 Lymphocytes mature in the bone marrow or the thymus and then congregate in lymphoid tissues throughout the body. 17
- 1-15 Adaptive immune responses are initiated by antigen and antigen-presenting cells in secondary lymphoid tissues. 18

- 1-16 Lymphocytes encounter and respond to antigen in the peripheral lymphoid organs. 19
- 1-17 Mucosal surfaces have specialized immune structures that orchestrate responses to environmental microbial encounters. 22
- 1-18 Lymphocytes activated by antigen proliferate in the peripheral lymphoid organs, generating effector cells and immunological memory. 23
- Summary. 24
- The effector mechanisms of immunity. 25**
- 1-19 Innate immune responses can select from several effector modules to protect against different types of pathogens. 26
- 1-20 Antibodies protect against extracellular pathogens and their toxic products. 27
- 1-21 T cells orchestrate cell-mediated immunity and regulate B-cell responses to most antigens. 29
- 1-22 Inherited and acquired defects in the immune system result in increased susceptibility to infection. 31
- 1-23 Understanding adaptive immune responses is important for the control of allergies, autoimmune disease, and the rejection of transplanted organs. 32
- 1-24 Vaccination is the most effective means of controlling infectious diseases. 33
- Summary. 34
- Summary to Chapter 1. 34
- Questions. 35
- References. 36

Chapter 2 Innate Immunity: The First Lines of Defense 37

- Anatomic barriers and initial chemical defenses. 38**
- 2-1 Infectious diseases are caused by diverse living agents that replicate in their hosts. 38
- 2-2 Epithelial surfaces of the body provide the first barrier against infection. 42
- 2-3 Infectious agents must overcome innate host defenses to establish a focus of infection. 44
- 2-4 Epithelial cells and phagocytes produce several kinds of antimicrobial proteins. 45
- Summary. 48
- The complement system and innate immunity. 49**
- 2-5 The complement system recognizes features of microbial surfaces and marks them for destruction by coating them with C3b. 50
- 2-6 The lectin pathway uses soluble receptors that recognize microbial surfaces to activate the complement cascade. 53

2-7	The classical pathway is initiated by activation of the C1 complex and is homologous to the lectin pathway.	56	3-8	The NOD-like receptors are intracellular sensors of bacterial infection and cellular damage.	96
2-8	Complement activation is largely confined to the surface on which it is initiated.	57	3-9	NLRP proteins react to infection or cellular damage through an inflammasome to induce cell death and inflammation.	98
2-9	The alternative pathway is an amplification loop for C3b formation that is accelerated by properdin in the presence of pathogens.	58	3-10	The RIG-I-like receptors detect cytoplasmic viral RNAs and activate MAVS to induce type I interferon production and pro-inflammatory cytokines.	101
2-10	Membrane and plasma proteins that regulate the formation and stability of C3 convertases determine the extent of complement activation.	60	3-11	Cytosolic DNA sensors signal through STING to induce production of type I interferons.	103
2-11	Complement developed early in the evolution of multicellular organisms.	61	3-12	Activation of innate sensors in macrophages and dendritic cells triggers changes in gene expression that have far-reaching effects on the immune response.	104
2-12	Surface-bound C3 convertase deposits large numbers of C3b fragments on pathogen surfaces and generates C5 convertase activity.	62	3-13	Toll signaling in <i>Drosophila</i> is downstream of a distinct set of pathogen-recognition molecules.	105
2-13	Ingestion of complement-tagged pathogens by phagocytes is mediated by receptors for the bound complement proteins.	63	3-14	TLR and NOD genes have undergone extensive diversification in both invertebrates and some primitive chordates.	106
2-14	The small fragments of some complement proteins initiate a local inflammatory response.	65		Summary.	106
2-15	The terminal complement proteins polymerize to form pores in membranes that can kill certain pathogens.	66		Induced innate responses to infection.	107
2-16	Complement control proteins regulate all three pathways of complement activation and protect the host from their destructive effects.	67	3-15	Cytokines and their receptors fall into distinct families of structurally related proteins.	107
2-17	Pathogens produce several types of proteins that can inhibit complement activation.	71	3-16	Cytokine receptors of the hematopoietin family are associated with the JAK family of tyrosine kinases, which activate STAT transcription factors.	109
	Summary.	72	3-17	Chemokines released by macrophages and dendritic cells recruit effector cells to sites of infection.	111
	Summary to Chapter 2.	73	3-18	Cell-adhesion molecules control interactions between leukocytes and endothelial cells during an inflammatory response.	113
	Questions.	74	3-19	Neutrophils make up the first wave of cells that cross the blood vessel wall to enter an inflamed tissue.	116
	References.	75	3-20	TNF- α is an important cytokine that triggers local containment of infection but induces shock when released systemically.	118
	Chapter 3 The Induced Responses of Innate Immunity	77	3-21	Cytokines made by macrophages and dendritic cells induce a systemic reaction known as the acute-phase response.	118
	Pattern recognition by cells of the innate immune system.	77	3-22	Interferons induced by viral infection make several contributions to host defense.	121
3-1	After entering tissues, many microbes are recognized, ingested, and killed by phagocytes.	78	3-23	Several types of innate lymphoid cells provide protection in early infection.	124
3-2	G-protein-coupled receptors on phagocytes link microbe recognition with increased efficiency of intracellular killing.	81	3-24	NK cells are activated by type I interferon and macrophage-derived cytokines.	125
3-3	Microbial recognition and tissue damage initiate an inflammatory response.	85	3-25	NK cells express activating and inhibitory receptors to distinguish between healthy and infected cells.	126
3-4	Toll-like receptors represent an ancient pathogen-recognition system.	87	3-26	NK-cell receptors belong to several structural families, the KIRs, KLRs, and NCRs.	128
3-5	Mammalian Toll-like receptors are activated by many different pathogen-associated molecular patterns.	88	3-27	NK cells express activating receptors that recognize ligands induced on infected cells or tumor cells.	130
3-6	TLR-4 recognizes bacterial lipopolysaccharide in association with the host accessory proteins MD-2 and CD14.	92		Summary.	131
3-7	TLRs activate NF κ B, AP-1, and IRF transcription factors to induce the expression of inflammatory cytokines and type I interferons.	92			

Summary to Chapter 3.	131	4-18	The CD4 and CD8 cell-surface proteins of T cells directly contact MHC molecules and are required to make an effective response to antigen.	163
Questions.	132			
References.	133	4-19	The two classes of MHC molecules are expressed differentially on cells.	166
PART II THE RECOGNITION OF ANTIGEN				
Chapter 4 Antigen Recognition by B-cell and T-cell Receptors				
The structure of a typical antibody molecule.	139	4-20	A distinct subset of T cells bears an alternative receptor made up of γ and δ chains.	166
4-1 IgG antibodies consist of four polypeptide chains.	141	Summary.		167
4-2 Immunoglobulin heavy and light chains are composed of constant and variable regions.	142	Summary to Chapter 4.		168
4-3 The domains of an immunoglobulin molecule have similar structures.	142	Questions.		169
4-4 The antibody molecule can readily be cleaved into functionally distinct fragments.	144	References.		170
4-5 The hinge region of the immunoglobulin molecule allows flexibility in binding to multiple antigens.	145	Chapter 5 The Generation of Lymphocyte Antigen Receptors		
Summary.	145	Primary immunoglobulin gene rearrangement.		
The interaction of the antibody molecule with specific antigen.	146	5-1	Immunoglobulin genes are rearranged in the progenitors of antibody-producing cells.	174
4-6 Localized regions of hypervariable sequence form the antigen-binding site.	146	5-2	Complete genes that encode a variable region are generated by the somatic recombination of separate gene segments.	175
4-7 Antibodies bind antigens via contacts in CDRs that are complementary to the size and shape of the antigen.	147	5-3	Multiple contiguous V gene segments are present at each immunoglobulin locus.	176
4-8 Antibodies bind to conformational shapes on the surfaces of antigens using a variety of noncovalent forces.	148	5-4	Rearrangement of V, D, and J gene segments is guided by flanking DNA sequences.	178
4-9 Antibody interaction with intact antigens is influenced by steric constraints.	150	5-5	The reaction that recombines V, D, and J gene segments involves both lymphocyte-specific and ubiquitous DNA-modifying enzymes.	179
4-10 Some species generate antibodies with alternative structures.	151	5-6	The diversity of the immunoglobulin repertoire is generated by four main processes.	184
Summary.	152	5-7	The multiple inherited gene segments are used in different combinations.	184
Antigen recognition by T cells.	152	5-8	Variable addition and subtraction of nucleotides at the junctions between gene segments contributes to the diversity of the third hypervariable region.	185
4-11 The TCR α : β heterodimer is very similar to a Fab fragment of immunoglobulin.	153	Summary.		186
4-12 A T-cell receptor recognizes antigen in the form of a complex of a foreign peptide bound to an MHC molecule.	155	T-cell receptor gene rearrangement.		
4-13 There are two classes of MHC molecules with distinct subunit compositions but similar three-dimensional structures.	155	5-9	The T-cell receptor gene segments are arranged in a similar pattern to immunoglobulin gene segments and are rearranged by the same enzymes.	187
4-14 Peptides are stably bound to MHC molecules, and also serve to stabilize the MHC molecule on the cell surface.	158	5-10	T-cell receptors concentrate diversity in the third hypervariable region.	189
4-15 MHC class I molecules bind short peptides of 8–10 amino acids by both ends.	158	5-11	γ : δ T-cell receptors are also generated by gene rearrangement.	190
4-16 The length of the peptides bound by MHC class II molecules is not constrained.	160	Summary.		191
4-17 The crystal structures of several peptide:MHC:T-cell receptor complexes show a similar orientation of the T-cell receptor over the peptide:MHC complex.	161	Structural variation in immunoglobulin constant regions.		
		5-12	Different classes of immunoglobulins are distinguished by the structure of their heavy-chain constant regions.	192
		5-13	The constant region confers functional specialization on the antibody.	193
		5-14	IgM and IgD are derived from the same pre-mRNA transcript and are both expressed on the surface of mature B cells.	194

5-15	Transmembrane and secreted forms of immunoglobulin are generated from alternative heavy-chain mRNA transcripts.	195	The major histocompatibility complex and its function.	231	
5-16	IgM and IgA can form polymers by interacting with the J chain.	197	6-10	Many proteins involved in antigen processing and presentation are encoded by genes within the MHC.	231
Summary.		198	6-11	The protein products of MHC class I and class II genes are highly polymorphic.	234
Evolution of the adaptive immune response.		198	6-12	MHC polymorphism affects antigen recognition by T cells by influencing both peptide binding and the contacts between T-cell receptor and MHC molecule.	235
5-17	Some invertebrates generate extensive diversity in a repertoire of immunoglobulin-like genes.	198	6-13	Alloreactive T cells recognizing nonself MHC molecules are very abundant.	239
5-18	Agnathans possess an adaptive immune system that uses somatic gene rearrangement to diversify receptors built from LRR domains.	200	6-14	Many T cells respond to superantigens.	240
5-19	RAG-dependent adaptive immunity based on a diversified repertoire of immunoglobulin-like genes appeared abruptly in the cartilaginous fishes.	202	6-15	MHC polymorphism extends the range of antigens to which the immune system can respond.	241
5-20	Different species generate immunoglobulin diversity in different ways.	203	Summary.		242
5-21	Both α : β and γ : δ T-cell receptors are present in cartilaginous fishes.	206	Generation of ligands for unconventional T-cell subsets.	242	
5-22	MHC class I and class II molecules are also first found in the cartilaginous fishes.	206	6-16	A variety of genes with specialized functions in immunity are also encoded in the MHC.	243
Summary.		207	6-17	Specialized MHC class I molecules act as ligands for the activation and inhibition of NK cells and unconventional T-cell subsets.	245
Summary to Chapter 5.		207	6-18	Members of the CD1 family of MHC class I-like molecules present microbial lipids to invariant NKT cells.	246
Questions.		208	6-19	The nonclassical MHC class I molecule MR1 presents microbial folate metabolites to MAIT cells.	248
References.		209	6-20	γ : δ T cells can recognize a variety of diverse ligands.	249
Chapter 6 Antigen Presentation to T Lymphocytes		213	Summary.		250
The generation of α:β T-cell receptor ligands.		214	Summary to Chapter 6.		250
6-1	Antigen presentation functions both in arming effector T cells and in triggering their effector functions to attack pathogen-infected cells.	214	Questions.		251
6-2	Peptides are generated from ubiquitinated proteins in the cytosol by the proteasome.	216	References.		252
6-3	Peptides from the cytosol are transported by TAP into the endoplasmic reticulum and further processed before binding to MHC class I molecules.	218	PART III THE DEVELOPMENT OF MATURE LYMPHOCYTE RECEPTOR REPERTOIRES		
6-4	Newly synthesized MHC class I molecules are retained in the endoplasmic reticulum until they bind a peptide.	219	Chapter 7 Lymphocyte Receptor Signaling	257	
6-5	Dendritic cells use cross-presentation to present exogenous proteins on MHC class I molecules to prime CD8 T cells.	222	General principles of signal transduction and propagation.	257	
6-6	Peptide:MHC class II complexes are generated in acidified endocytic vesicles from proteins obtained through endocytosis, phagocytosis, and autophagy.	223	7-1	Transmembrane receptors convert extracellular signals into intracellular biochemical events.	258
6-7	The invariant chain directs newly synthesized MHC class II molecules to acidified intracellular vesicles.	225	7-2	Intracellular signal propagation is mediated by large multiprotein signaling complexes.	260
6-8	The MHC class II-like molecules HLA-DM and HLA-DO regulate exchange of CLIP for other peptides.	226	7-3	Small G proteins act as molecular switches in many different signaling pathways.	262
6-9	Cessation of antigen processing occurs in dendritic cells after their activation through reduced expression of the MARCH-1 E3 ligase.	229	7-4	Signaling proteins are recruited to the membrane by a variety of mechanisms.	262
Summary.		230	7-5	Post-translational modifications of proteins can both activate and inhibit signaling responses.	263
			7-6	The activation of some receptors generates small-molecule second messengers.	264
			Summary.		265

Antigen receptor signaling and lymphocyte activation.	265	Questions.	290
7-7 Antigen receptors consist of variable antigen-binding chains associated with invariant chains that carry out the signaling function of the receptor.	266	References.	291
7-8 Antigen recognition by the T-cell receptor and its co-receptors transduces a signal across the plasma membrane to initiate signaling.	267	Chapter 8 The Development of B and T Lymphocytes	295
7-9 Antigen recognition by the T-cell receptor and its co-receptors leads to phosphorylation of ITAMs by Src-family kinases, generating the first intracellular signal in a signaling cascade.	268	Development of B lymphocytes.	296
7-10 Phosphorylated ITAMs recruit and activate the tyrosine kinase ZAP-70.	270	8-1 Lymphocytes derive from hematopoietic stem cells in the bone marrow.	297
7-11 ITAMs are also found in other receptors on leukocytes that signal for cell activation.	270	8-2 B-cell development begins by rearrangement of the heavy-chain locus.	299
7-12 Activated ZAP-70 phosphorylates scaffold proteins and promotes PI 3-kinase activation.	271	8-3 The pre-B-cell receptor tests for successful production of a complete heavy chain and signals for the transition from the pro-B cell to the pre-B cell stage.	302
7-13 Activated PLC- γ generates the second messengers diacylglycerol and inositol trisphosphate that lead to transcription factor activation.	272	8-4 Pre-B-cell receptor signaling inhibits further heavy-chain locus rearrangement and enforces allelic exclusion.	303
7-14 Ca ²⁺ entry activates the transcription factor NFAT.	273	8-5 Pre-B cells rearrange the light-chain locus and express cell-surface immunoglobulin.	304
7-15 Ras activation stimulates the mitogen-activated protein kinase (MAPK) relay and induces expression of the transcription factor AP-1.	274	8-6 Immature B cells are tested for autoreactivity before they leave the bone marrow.	305
7-16 Protein kinase C activates the transcription factors NF κ B and AP-1.	276	8-7 Lymphocytes that encounter sufficient quantities of self antigens for the first time in the periphery are eliminated or inactivated.	308
7-17 PI 3-kinase activation upregulates cellular metabolic pathways via the serine/threonine kinase Akt.	277	8-8 Immature B cells arriving in the spleen turn over rapidly and require cytokines and positive signals through the B-cell receptor for maturation and long-term survival.	309
7-18 T-cell receptor signaling leads to enhanced integrin-mediated cell adhesion.	278	8-9 B-1 B cells are an innate lymphocyte subset that arises early in development.	312
7-19 T-cell receptor signaling induces cytoskeletal reorganization by activating the small GTPase Cdc42.	279	Summary.	313
7-20 The logic of B-cell receptor signaling is similar to that of T-cell receptor signaling, but some of the signaling components are specific to B cells.	279	Development of T lymphocytes.	315
Summary.	282	8-10 T-cell progenitors originate in the bone marrow, but all the important events in their development occur in the thymus.	315
Co-stimulatory and inhibitory receptors modulate antigen receptor signaling in T and B lymphocytes.	282	8-11 Commitment to the T-cell lineage occurs in the thymus following Notch signaling.	317
7-21 The cell-surface protein CD28 is a required co-stimulatory signaling receptor for naive T-cell activation.	283	8-12 T-cell precursors proliferate extensively in the thymus, but most die there.	317
7-22 Maximal activation of PLC- γ , which is important for transcription factor activation, requires a co-stimulatory signal induced by CD28.	284	8-13 Successive stages in the development of thymocytes are marked by changes in cell-surface molecules.	319
7-23 TNF receptor superfamily members augment T-cell and B-cell activation.	284	8-14 Thymocytes at different developmental stages are found in distinct parts of the thymus.	321
7-24 Inhibitory receptors on lymphocytes downregulate immune responses by interfering with co-stimulatory signaling pathways.	286	8-15 T cells with α : β or γ : δ receptors arise from a common progenitor.	322
7-25 Inhibitory receptors on lymphocytes downregulate immune responses by recruiting protein or lipid phosphatases.	287	8-16 T cells expressing γ : δ T-cell receptors arise in two distinct phases during development.	322
Summary.	288	8-17 Successful synthesis of a rearranged β chain allows the production of a pre-T-cell receptor that triggers cell proliferation and blocks further β -chain gene rearrangement.	324
Summary to Chapter 7.	289	8-18 T-cell α -chain genes undergo successive rearrangements until positive selection or cell death intervenes.	326
		Summary.	328

Positive and negative selection of T cells.	328	9-9	Dendritic cells process antigens from a wide array of pathogens.	358
8-19 Only thymocytes whose receptors interact with self peptide:self MHC complexes can survive and mature.	328	9-10	Microbe-induced TLR signaling in tissue-resident dendritic cells induces their migration to lymphoid organs and enhances antigen processing.	361
8-20 Positive selection acts on a repertoire of T-cell receptors with inherent specificity for MHC molecules.	329	9-11	Plasmacytoid dendritic cells produce abundant type I interferons and may act as helper cells for antigen presentation by conventional dendritic cells.	363
8-21 Positive selection coordinates the expression of CD4 or CD8 with the specificity of the T-cell receptor and the potential effector functions of the T cell.	330	9-12	Macrophages are scavenger cells that can be induced by pathogens to present foreign antigens to naive T cells.	363
8-22 Thymic cortical epithelial cells mediate positive selection of developing thymocytes.	331	9-13	B cells are highly efficient at presenting antigens that bind to their surface immunoglobulin.	364
8-23 T cells that react strongly with ubiquitous self antigens are deleted in the thymus.	332		Summary.	366
8-24 Negative selection is driven most efficiently by bone marrow-derived antigen-presenting cells.	334		Priming of naive T cells by pathogen-activated dendritic cells.	366
8-25 The specificity and/or the strength of signals for negative and positive selection must differ.	334	9-14	Cell-adhesion molecules mediate the initial interaction of naive T cells with antigen-presenting cells.	367
8-26 Self-recognizing regulatory T cells and innate T cells develop in the thymus.	335	9-15	Antigen-presenting cells deliver multiple signals for the clonal expansion and differentiation of naive T cells.	368
8-27 The final stage of T-cell maturation occurs in the thymic medulla.	336	9-16	CD28-dependent co-stimulation of activated T cells induces expression of interleukin-2 and the high-affinity IL-2 receptor.	368
8-28 T cells that encounter sufficient quantities of self antigens for the first time in the periphery are eliminated or inactivated.	336	9-17	Additional co-stimulatory pathways are involved in T-cell activation.	369
Summary.	337	9-18	Proliferating T cells differentiate into effector T cells that do not require co-stimulation to act.	370
Summary to Chapter 8.	337	9-19	CD8 T cells can be activated in different ways to become cytotoxic effector cells.	372
Questions.	339	9-20	CD4 T cells differentiate into several subsets of functionally different effector cells.	372
References.	340	9-21	Cytokines induce the differentiation of naive CD4 T cells down distinct effector pathways.	375
		9-22	CD4 T-cell subsets can cross-regulate each other's differentiation through the cytokines they produce.	377
		9-23	Regulatory CD4 T cells are involved in controlling adaptive immune responses.	379
			Summary.	380
PART IV THE ADAPTIVE IMMUNE RESPONSE	345		General properties of effector T cells and their cytokines.	380
Chapter 9 T-cell-Mediated Immunity	345	9-24	Effector T-cell interactions with target cells are initiated by antigen-nonspecific cell-adhesion molecules.	381
Development and function of secondary lymphoid organs—sites for the initiation of adaptive immune responses.	347	9-25	An immunological synapse forms between effector T cells and their targets to regulate signaling and to direct the release of effector molecules.	381
9-1 T and B lymphocytes are found in distinct locations in secondary lymphoid tissues.	347	9-26	The effector functions of T cells are determined by the array of effector molecules that they produce.	383
9-2 The development of secondary lymphoid tissues is controlled by lymphoid tissue inducer cells and proteins of the tumor necrosis factor family.	349	9-27	Cytokines can act locally or at a distance.	383
9-3 T and B cells are partitioned into distinct regions of secondary lymphoid tissues by the actions of chemokines.	350	9-28	T cells express several TNF-family cytokines as trimeric proteins that are usually associated with the cell surface.	386
9-4 Naive T cells migrate through secondary lymphoid tissues, sampling peptide:MHC complexes on dendritic cells.	351		Summary.	386
9-5 Lymphocyte entry into lymphoid tissues depends on chemokines and adhesion molecules.	352			
9-6 Activation of integrins by chemokines is responsible for the entry of naive T cells into lymph nodes.	353			
9-7 The exit of T cells from lymph nodes is controlled by a chemotactic lipid.	355			
9-8 T-cell responses are initiated in secondary lymphoid organs by activated dendritic cells.	356			

T-cell-mediated cytotoxicity.	387	10-14 Some antigens do not require T-cell help to induce B-cell responses.	419
9-29 Cytotoxic T cells induce target cells to undergo programmed cell death via extrinsic and intrinsic pathways of apoptosis.	387	Summary.	421
9-30 The intrinsic pathway of apoptosis is mediated by the release of cytochrome c from mitochondria.	389	The distributions and functions of immunoglobulin classes.	422
9-31 Cytotoxic effector proteins that trigger apoptosis are contained in the granules of CD8 cytotoxic T cells.	390	10-15 Antibodies of different classes operate in distinct places and have distinct effector functions.	423
9-32 Cytotoxic T cells are selective serial killers of targets expressing a specific antigen.	391	10-16 Polymeric immunoglobulin receptor binds to the Fc regions of IgA and IgM and transports them across epithelial barriers.	425
9-33 Cytotoxic T cells also act by releasing cytokines.	392	10-17 The neonatal Fc receptor carries IgG across the placenta and prevents IgG excretion from the body.	426
Summary.	392	10-18 High-affinity IgG and IgA antibodies can neutralize toxins and block the infectivity of viruses and bacteria.	426
Summary to Chapter 9.	392	10-19 Antibody:antigen complexes activate the classical pathway of complement by binding to C1q.	429
Questions.	393	10-20 Complement receptors and Fc receptors both contribute to removal of immune complexes from the circulation.	430
References.	395	Summary.	431
Chapter 10 The Humoral Immune Response	399	The destruction of antibody-coated pathogens via Fc receptors.	432
B-cell activation by antigen and helper T cells.	400	10-21 The Fc receptors of accessory cells are signaling receptors specific for immunoglobulins of different classes.	432
10-1 Activation of B cells by antigen involves signals from the B-cell receptor and either T _{FH} cells or microbial antigens.	400	10-22 Fc receptors on phagocytes are activated by antibodies bound to the surface of pathogens and enable the phagocytes to ingest and destroy pathogens.	433
10-2 Linked recognition of antigen by T cells and B cells promotes robust antibody responses.	402	10-23 Fc receptors activate NK cells to destroy antibody-coated targets.	435
10-3 B cells that encounter their antigens migrate toward the boundaries between B-cell and T-cell areas in secondary lymphoid tissues.	403	10-24 Mast cells and basophils bind IgE antibody via the high-affinity Fcε receptor.	436
10-4 T cells express surface molecules and cytokines that activate B cells, which in turn promote T _{FH} -cell development.	406	10-25 IgE-mediated activation of accessory cells has an important role in resistance to parasite infection.	437
10-5 Activated B cells differentiate into antibody-secreting plasmablasts and plasma cells.	406	Summary.	438
10-6 The second phase of a primary B-cell immune response occurs when activated B cells migrate into follicles and proliferate to form germinal centers.	408	Summary to Chapter 10.	439
10-7 Germinal center B cells undergo V-region somatic hypermutation, and cells with mutations that improve affinity for antigen are selected.	410	Questions.	440
10-8 Positive selection of germinal center B cells involves contact with T _{FH} cells and CD40 signaling.	412	References.	441
10-9 Activation-induced cytidine deaminase (AID) introduces mutations into genes transcribed in B cells.	413	Chapter 11 Integrated Dynamics of Innate and Adaptive Immunity	445
10-10 Mismatch and base-excision repair pathways contribute to somatic hypermutation following initiation by AID.	414	Integration of innate and adaptive immunity in response to specific types of pathogens.	446
10-11 AID initiates class switching to allow the same assembled V _H exon to be associated with different C _H genes in the course of an immune response.	415	11-1 The course of an infection can be divided into several distinct phases.	446
10-12 Cytokines made by T _{FH} cells direct the choice of isotype for class switching in T-dependent antibody responses.	418	11-2 The effector mechanisms that are recruited to clear an infection depend on the infectious agent.	449
10-13 B cells that survive the germinal center reaction eventually differentiate into either plasma cells or memory cells.	419	Summary.	452
		Effector T cells augment the effector functions of innate immune cells.	452
		11-3 Effector T cells are guided to specific tissues and sites of infection by changes in their expression of adhesion molecules and chemokine receptors.	453

11-4	Pathogen-specific effector T cells are enriched at sites of infection as adaptive immunity progresses.	457	Summary.	485
			Summary to Chapter 11.	486
11-5	T_H1 cells coordinate and amplify the host response to intracellular pathogens through classical activation of macrophages.	458	Questions.	487
			References.	488
11-6	Activation of macrophages by T_H1 cells must be tightly regulated to avoid tissue damage.	460	Chapter 12 The Mucosal Immune System	493
11-7	Chronic activation of macrophages by T_H1 cells mediates the formation of granulomas to contain intracellular pathogens that cannot be cleared.	461	The nature and structure of the mucosal immune system.	493
11-8	Defects in type 1 immunity reveal its important role in the elimination of intracellular pathogens.	461	12-1 The mucosal immune system protects the internal surfaces of the body.	493
11-9	T_H2 cells coordinate type 2 responses to expel intestinal helminths and repair tissue injury.	462	12-2 Cells of the mucosal immune system are located both in anatomically defined compartments and scattered throughout mucosal tissues.	496
11-10	T_H17 cells coordinate type 3 responses to enhance the clearance of extracellular bacteria and fungi.	465	12-3 The intestine has distinctive routes and mechanisms of antigen uptake.	499
11-11	Differentiated effector T cells continue to respond to signals as they carry out their effector functions.	466	12-4 The mucosal immune system contains large numbers of effector lymphocytes even in the absence of disease.	500
11-12	Effector T cells can be activated to release cytokines independently of antigen recognition.	467	12-5 The circulation of lymphocytes within the mucosal immune system is controlled by tissue-specific adhesion molecules and chemokine receptors.	501
11-13	Effector T cells demonstrate plasticity and cooperativity that enable adaptation during anti-pathogen responses.	468	12-6 Priming of lymphocytes in one mucosal tissue may induce protective immunity at other mucosal surfaces.	502
11-14	Integration of cell- and antibody-mediated immunity is critical for protection against many types of pathogens.	469	12-7 Distinct populations of dendritic cells control mucosal immune responses.	503
11-15	Primary CD8 T-cell responses to pathogens can occur in the absence of CD4 T-cell help.	470	12-8 Macrophages and dendritic cells have different roles in mucosal immune responses.	505
11-16	Resolution of an infection is accompanied by the death of most of the effector cells and the generation of memory cells.	471	12-9 Antigen-presenting cells in the intestinal mucosa acquire antigen by a variety of routes.	505
	Summary.	472	12-10 Secretory IgA is the class of antibody associated with the mucosal immune system.	506
	Immunological memory.	473	12-11 T-independent processes can contribute to IgA production in some species.	509
11-17	Immunological memory is long lived after infection or vaccination.	473	12-12 IgA deficiency is relatively common in humans but may be compensated for by secretory IgM.	509
11-18	Memory B-cell responses are more rapid and have higher affinity for antigen compared with responses of naive B cells.	475	12-13 The intestinal lamina propria contains antigen-experienced T cells and populations of unusual innate lymphoid cells.	510
11-19	Memory B cells can reenter germinal centers and undergo additional somatic hypermutation and affinity maturation during secondary immune responses.	476	12-14 The intestinal epithelium is a unique compartment of the immune system.	511
11-20	MHC tetramers identify memory T cells that persist at an increased frequency relative to their frequency as naive T cells.	477	Summary.	514
11-21	Memory T cells arise from effector T cells that maintain sensitivity to IL-7 or IL-15.	478	The mucosal response to infection and regulation of mucosal immune responses.	514
11-22	Memory T cells are heterogeneous and include central memory, effector memory, and tissue-resident subsets.	480	12-15 Enteric pathogens cause a local inflammatory response and the development of protective immunity.	515
11-23	CD4 T-cell help is required for CD8 T-cell memory and involves CD40 and IL-2 signaling.	482	12-16 Pathogens induce adaptive immune responses when innate defenses have been breached.	518
11-24	In immune individuals, secondary and subsequent responses are mainly attributable to memory lymphocytes.	484	12-17 Effector T-cell responses in the intestine protect the function of the epithelium.	518
			12-18 The mucosal immune system must maintain tolerance to harmless foreign antigens.	519
			12-19 The normal intestine contains large quantities of bacteria that are required for health.	520

12-20	Innate and adaptive immune systems control microbiota while preventing inflammation without compromising the ability to react to invaders.	521	13-16	Mutations in the molecular regulators of inflammation can cause uncontrolled inflammatory responses that result in 'autoinflammatory disease.'	556
12-21	The intestinal microbiota plays a major role in shaping intestinal and systemic immune function.	522	13-17	Hematopoietic stem cell transplantation or gene therapy can be useful to correct genetic defects.	557
12-22	Full immune responses to commensal bacteria provoke intestinal disease.	524	13-18	Noninherited, secondary immunodeficiencies are major predisposing causes of infection and death.	558
Summary.		525	Summary.		559
Summary to Chapter 12.		525	Evasion and subversion of immune defenses.		560
Questions.		526	13-19	Extracellular bacterial pathogens have evolved different strategies to avoid detection by pattern recognition receptors and destruction by antibody, complement, and antimicrobial peptides.	560
References.		527	13-20	Intracellular bacterial pathogens can evade the immune system by seeking shelter within phagocytes.	563

PART V THE IMMUNE SYSTEM IN HEALTH AND DISEASE

Chapter 13 Failures of Host Defense Mechanisms 533

Immunodeficiency diseases. 533

13-1	A history of repeated infections suggests a diagnosis of immunodeficiency.	534	13-22	RNA viruses use different mechanisms of antigenic variation to keep a step ahead of the adaptive immune system.	566
13-2	Primary immunodeficiency diseases are caused by inherited gene defects.	534	13-23	DNA viruses use multiple mechanisms to subvert NK-cell and CTL responses.	568
13-3	Defects in T-cell development can result in severe combined immunodeficiencies.	535	13-24	Some latent viruses persist <i>in vivo</i> by ceasing to replicate until immunity wanes.	571
13-4	SCID can also be due to defects in the purine salvage pathway.	538	Summary.		573
13-5	Defects in antigen receptor gene rearrangement can result in SCID.	538	Acquired immune deficiency syndrome.		573
13-6	Defects in signaling from T-cell antigen receptors can cause severe immunodeficiency.	539	13-25	HIV is a retrovirus that establishes a chronic infection that slowly progresses to AIDS.	574
13-7	Genetic defects in thymic function that block T-cell development result in severe immunodeficiencies.	539	13-26	HIV infects and replicates within cells of the immune system.	576
13-8	Defects in B-cell development result in deficiencies in antibody production that cause an inability to clear extracellular bacteria and some viruses.	541	13-27	Activated CD4 T cells are the major source of HIV replication.	578
13-9	Immune deficiencies can be caused by defects in B-cell or T-cell activation and function that lead to abnormal antibody responses.	543	13-28	There are several routes by which HIV is transmitted and establishes infection.	579
13-10	Normal pathways for host defense against different infectious agents are pinpointed by genetic deficiencies of cytokine pathways central to type 1/T _H 1 and type 3/T _H 17 responses.	546	13-29	HIV variants with tropism for different co-receptors play different roles in transmission and progression of disease.	580
13-11	Inherited defects in the cytolytic pathway of lymphocytes can cause uncontrolled lymphoproliferation and inflammatory responses to viral infections.	548	13-30	A genetic deficiency of the co-receptor CCR5 confers resistance to HIV infection.	582
13-12	X-linked lymphoproliferative syndrome is associated with fatal infection by Epstein–Barr virus and with the development of lymphomas.	550	13-31	An immune response controls but does not eliminate HIV.	583
13-13	Immunodeficiency is caused by inherited defects in the development of dendritic cells.	551	13-32	Lymphoid tissue is the major reservoir of HIV infection.	585
13-14	Defects in complement components and complement-regulatory proteins cause defective humoral immune function and tissue damage.	552	13-33	Genetic variation in the host can alter the rate of disease progression.	585
13-15	Defects in phagocytic cells permit widespread bacterial infections.	553	13-34	The destruction of immune function as a result of HIV infection leads to increased susceptibility to opportunistic infection and eventually to death.	587
			13-35	Drugs that block HIV replication lead to a rapid decrease in titer of infectious virus and an increase in CD4 T cells.	588
			13-36	In the course of infection HIV accumulates many mutations, which can result in the outgrowth of drug-resistant variants.	590

13-37	Vaccination against HIV is an attractive solution but poses many difficulties.	591	14-17	Celiac disease has features of both an allergic response and autoimmunity.	634
13-38	Prevention and education are important in controlling the spread of HIV and AIDS.	592		Summary.	636
	Summary.	593		Summary to Chapter 14.	636
	Summary to Chapter 13.	594		Questions.	637
	Questions.	594		References.	638
	References.	595		Chapter 15 Autoimmunity and Transplantation	643
	Chapter 14 Allergy and Allergic Diseases	601		The making and breaking of self-tolerance.	643
	IgE and IgE-mediated allergic diseases.	602	15-1	A critical function of the immune system is to discriminate self from nonself.	643
14-1	Sensitization involves class switching to IgE production on first contact with an allergen.	603	15-2	Multiple tolerance mechanisms normally prevent autoimmunity.	645
14-2	Although many types of antigens can cause allergic sensitization, proteases are common sensitizing agents.	605	15-3	Central deletion or inactivation of newly formed lymphocytes is the first checkpoint of self-tolerance.	646
14-3	Genetic factors contribute to the development of IgE-mediated allergic disease.	607	15-4	Lymphocytes that bind self antigens with relatively low affinity usually ignore them but in some circumstances become activated.	647
14-4	Environmental factors may interact with genetic susceptibility to cause allergic disease.	609	15-5	Antigens in immunologically privileged sites do not induce immune attack but can serve as targets.	648
14-5	Regulatory T cells can control allergic responses.	611	15-6	Autoreactive T cells that express particular cytokines may be nonpathogenic or may suppress pathogenic lymphocytes.	649
	Summary.	612	15-7	Autoimmune responses can be controlled at various stages by regulatory T cells.	650
	Effector mechanisms in IgE-mediated allergic reactions.	612		Summary.	652
14-6	Most IgE is cell-bound and engages effector mechanisms of the immune system by pathways different from those of other antibody isotypes.	613		Autoimmune diseases and pathogenic mechanisms.	652
14-7	Mast cells reside in tissues and orchestrate allergic reactions.	613	15-8	Specific adaptive immune responses to self antigens can cause autoimmune disease.	652
14-8	Eosinophils and basophils cause inflammation and tissue damage in allergic reactions.	616	15-9	Autoimmunity can be classified into either organ-specific or systemic disease.	653
14-9	IgE-mediated allergic reactions have a rapid onset but can also lead to chronic responses.	617	15-10	Multiple components of the immune system are typically recruited in autoimmune disease.	654
14-10	Allergen introduced into the bloodstream can cause anaphylaxis.	619	15-11	Chronic autoimmune disease develops through positive feedback from inflammation, inability to clear the self antigen, and a broadening of the autoimmune response.	657
14-11	Allergen inhalation is associated with the development of rhinitis and asthma.	621	15-12	Both antibody and effector T cells can cause tissue damage in autoimmune disease.	659
14-12	Allergy to particular foods causes systemic reactions as well as symptoms limited to the gut.	624	15-13	Autoantibodies against blood cells promote their destruction.	661
14-13	IgE-mediated allergic disease can be treated by inhibiting the effector pathways that lead to symptoms or by desensitization techniques that aim at restoring biological tolerance to the allergen.	625	15-14	The fixation of sublytic doses of complement to cells in tissues stimulates a powerful inflammatory response.	661
	Summary.	627	15-15	Autoantibodies against receptors cause disease by stimulating or blocking receptor function.	662
	Non-IgE-mediated allergic diseases.	628	15-16	Autoantibodies against extracellular antigens cause inflammatory injury.	663
14-14	Non-IgE dependent drug-induced hypersensitivity reactions in susceptible individuals occur by binding of the drug to the surface of circulating blood cells.	628	15-17	T cells specific for self antigens can cause direct tissue injury and sustain autoantibody responses.	665
14-15	Systemic disease caused by immune-complex formation can follow the administration of large quantities of poorly catabolized antigens.	628		Summary.	668
14-16	Hypersensitivity reactions can be mediated by T _H 1 cells and CD8 cytotoxic T cells.	630		The genetic and environmental basis of autoimmunity.	669
			15-18	Autoimmune diseases have a strong genetic component.	669

15-19	Genomics-based approaches are providing new insight into the immunogenetic basis of autoimmunity.	670	16-2	Cytotoxic drugs cause immunosuppression by killing dividing cells and have serious side-effects.	703
15-20	Many genes that predispose to autoimmunity fall into categories that affect one or more tolerance mechanisms.	674	16-3	Cyclosporin A, tacrolimus, rapamycin, and JAK inhibitors are effective immunosuppressive agents that interfere with various T-cell signaling pathways.	704
15-21	Monogenic defects of immune tolerance.	674	16-4	Antibodies against cell-surface molecules can be used to eliminate lymphocyte subsets or to inhibit lymphocyte function.	706
15-22	MHC genes have an important role in controlling susceptibility to autoimmune disease.	676	16-5	Antibodies can be engineered to reduce their immunogenicity in humans.	707
15-23	Genetic variants that impair innate immune responses can predispose to T-cell-mediated chronic inflammatory disease.	678	16-6	Monoclonal antibodies can be used to prevent allograft rejection.	708
15-24	External events can initiate autoimmunity.	679	16-7	Depletion of autoreactive lymphocytes can treat autoimmune disease.	710
15-25	Infection can lead to autoimmune disease by providing an environment that promotes lymphocyte activation.	680	16-8	Biologics that block TNF- α , IL-1, or IL-6 can alleviate autoimmune diseases.	711
15-26	Cross-reactivity between foreign molecules on pathogens and self molecules can lead to antiseif responses and autoimmune disease.	680	16-9	Biologic agents can block cell migration to sites of inflammation and reduce immune responses.	712
15-27	Drugs and toxins can cause autoimmune syndromes.	682	16-10	Blockade of co-stimulatory pathways that activate lymphocytes can be used to treat autoimmune disease.	713
15-28	Random events may be required for the initiation of autoimmunity.	682	16-11	Some commonly used drugs have immunomodulatory properties.	713
Summary.		682	16-12	Controlled administration of antigen can be used to manipulate the nature of an antigen-specific response.	714
Responses to alloantigens and transplant rejection.		683	Summary.		714
15-29	Graft rejection is an immunological response mediated primarily by T cells.	683	Using the immune response to attack tumors.		716
15-30	Transplant rejection is caused primarily by the strong immune response to nonself MHC molecules.	684	16-13	The development of transplantable tumors in mice led to the discovery of protective immune responses to tumors.	716
15-31	In MHC-identical grafts, rejection is caused by peptides from other alloantigens bound to graft MHC molecules.	685	16-14	Tumors are 'edited' by the immune system as they evolve and can escape rejection in many ways.	717
15-32	There are two ways of presenting alloantigens on the transplanted donor organ to the recipient's T lymphocytes.	686	16-15	Tumor rejection antigens can be recognized by T cells and form the basis of immunotherapies.	720
15-33	Antibodies that react with endothelium cause hyperacute graft rejection.	688	16-16	T cells expressing chimeric antigen receptors are an effective treatment in some leukemias.	723
15-34	Late failure of transplanted organs is caused by chronic injury to the graft.	688	16-17	Monoclonal antibodies against tumor antigens, alone or linked to toxins, can control tumor growth.	724
15-35	A variety of organs are transplanted routinely in clinical medicine.	689	16-18	Enhancing the immune response to tumors by vaccination holds promise for cancer prevention and therapy.	726
15-36	The converse of graft rejection is graft-versus-host disease.	691	16-19	Checkpoint blockade can augment immune responses to existing tumors.	727
15-37	Regulatory T cells are involved in alloreactive immune responses.	692	Summary.		728
15-38	The fetus is an allograft that is tolerated repeatedly.	693	Fighting infectious diseases with vaccination.		729
Summary.		694	16-20	Vaccines can be based on attenuated pathogens or material from killed organisms.	730
Summary to Chapter 15.		694	16-21	Most effective vaccines generate antibodies that prevent the damage caused by toxins or that neutralize the pathogen and stop infection.	731
Questions.		695	16-22	Effective vaccines must induce long-lasting protection while being safe and inexpensive.	732
References.		696			
Chapter 16 Manipulation of the Immune Response		701			
Treatment of unwanted immune responses.		701			
16-1	Corticosteroids are powerful anti-inflammatory drugs that alter the transcription of many genes.	702			

16-23	Live-attenuated viral vaccines are usually more potent than 'killed' vaccines and can be made safer by the use of recombinant DNA technology.	732	A-16	Isolation of peripheral blood lymphocytes by density-gradient fractionation.	766	
16-24	Live-attenuated vaccines can be developed by selecting nonpathogenic or disabled bacteria or by creating genetically attenuated parasites (GAPs).	734	A-17	Isolation of lymphocytes from tissues other than blood.	766	
16-25	The route of vaccination is an important determinant of success.	735	A-18	Flow cytometry and FACS analysis.	767	
16-26	<i>Bordetella pertussis</i> vaccination illustrates the importance of the perceived safety of a vaccine.	736	A-19	Lymphocyte isolation using antibody-coated magnetic beads.	770	
16-27	Conjugate vaccines have been developed as a result of linked recognition between T and B cells.	737	A-20	Isolation of homogeneous T-cell lines.	770	
16-28	Peptide-based vaccines can elicit protective immunity, but they require adjuvants and must be targeted to the appropriate cells and cell compartment to be effective.	738	A-21	Limiting-dilution culture.	771	
16-29	Adjuvants are important for enhancing the immunogenicity of vaccines, but few are approved for use in humans.	739	A-22	ELISPOT assay.	773	
16-30	Protective immunity can be induced by DNA-based vaccination.	740	A-23	Identification of functional subsets of T cells based on cytokine production or transcription factor expression.	773	
16-31	Vaccination and checkpoint blockade may be useful in controlling existing chronic infections.	741	A-24	Identification of T-cell receptor specificity using peptide:MHC tetramers.	776	
Summary.		742	A-25	Biosensor assays for measuring the rates of association and dissociation of antigen receptors for their ligands.	777	
Summary to Chapter 16.		742	A-26	Assays of lymphocyte proliferation.	778	
Questions.		743	A-27	Measurements of apoptosis.	779	
References.		744	A-28	Assays for cytotoxic T cells.	780	
APPENDICES			A-29	Assays for CD4 T cells.	782	
Appendix I The Immunologist's Toolbox			749	A-30	Transfer of protective immunity.	782
A-1.	Immunization.	749	A-31	Adoptive transfer of lymphocytes.	783	
A-2	Antibody responses.	752	A-32	Hematopoietic stem-cell transfers.	784	
A-3	Affinity chromatography.	753	A-33	<i>In vivo</i> administration of antibodies.	785	
A-4	Radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), and competitive inhibition assay.	753	A-34	Transgenic mice.	786	
A-5	Hemagglutination and blood typing.	755	A-35	Gene knockout by targeted disruption.	786	
A-6	Coombs tests and the detection of rhesus incompatibility.	756	A-36	Knockdown of gene expression by RNA interference (RNAi).	790	
A-7	Monoclonal antibodies.	757	Appendix II CD antigens		791	
A-8	Phage display libraries for antibody V-region production.	758	Appendix II Cytokines and their Receptors		811	
A-9	Generation of human monoclonal antibodies from vaccinated individuals.	759	Appendix IV Chemokines and their Receptors		814	
A-10	Microscopy and imaging using fluorescent dyes.	760	Biographies		816	
A-11	Immunoelectron microscopy.	761	Photograph Acknowledgments		817	
A-12	Immunohistochemistry.	762	Glossary		818	
A-13	Immunoprecipitation and co-immunoprecipitation.	762	Index		855	
A-14	Immunoblotting (Western blotting).	764				
A-15	Use of antibodies in the isolation and characterization of multiprotein complexes by mass spectrometry.	764				